

Gravitational-Wave Bursts Induced by Neutrino Oscillations: The Origin of Asymmetry in Supernovae Explosions

Herman J. Mosquera Cuesta
High Energy/Astrophysics Section
Abdus Salam International Centre for Theoretical Physics
Strada Costiera 11, Miramare 34014, Trieste, Italy
e-mail: herman@ictp.trieste.it

(February 1, 2008)

If neutrino flavor changes really exist, to say: μ -neutrino oscillating into a *sterile* neutrino, then, it can be expected that due to neutrino oscillations and non-spherical distortion of the resonance surface induced by the magnetic field, some asymmetric emission of sterile neutrinos can occur during the protoneutron star formation at the onset of a supernova core-collapse. Assuming no strong suppression of the oscillations, the non-spherical huge neutrino energies released, ($\sim 10^{53-54}$ *erg*), together with the proto-neutron star rapid rotation, may trigger powerful bursts of gravitational waves by the time neutrino flavor conversions ensue. I show here that these bursts are detectable by the new generation of gravitational-wave detectors as LIGO, VIRGO and TIGAs for distance scales ~ 10 *kpc*. It is also argued that the general relativity requirement of an ellipsoidal axisymmetric core at maximum gravitational-wave emission induced by ν -luminosity can properly be afforded by the neutrino-sphere geometry when the oscillations onset. The connection of neutrino oscillations with the supernova asymmetry and bi-polar jets ejecta is shown naturally to appear in this scenario.

PACS numbers: 04.30.Db, 04.40.Dg, 14.60.Pq, 97.60.Bw

Turning the millenium a new generation of interferometric and resonant-mass gravitational-wave antennas will be operational. If confirmed, the detection of gravitational radiation will make a breakthrough in contemporary astrophysics and cosmology [1,2]. Due to the fundamental impact these observations would have, investigation of potential detectable sources becomes a key piece. In this *letter* I propound for first time that a new fundamental source of gravitational radiation had been overlooked by the community of workers on the fields. I suggest that the recently confirmed resonant conversion of neutrinos in dense magnetized matter can be a very important new one. I conjecture that this neutrino temperature asymmetry, produced during $\nu_\mu \leftrightarrow \nu_{\tau,s}$ flavor changes, couples to a non-spherical mass-energy distribution (the oscillating neutrino fluid) inside the proto-neutron star during a core-collapse supernova explosion. The resonant neutrino conversion and rapid rotation may provide the key ingredient for the analogous of a quadrupolar mass-moment tensor for the whole NS to develop, as currently is pictured in modeling GWs from nonaxisymmetric spinning neutron stars. Consequently, it makes it possible the generation of gravitational-wave bursts induced by neutrino conversions, $\nu_\mu \leftrightarrow \nu_{\tau,s}$, powered by the onset of the collapse of the inner core during explosions of massive stars.

In this *letter* it is argued that the neutrino oscillation process as described by Kusenko and Segrè [3] (see below), and its associated gravitational-wave burst, as suggested for the first time here, occurs during the very

early stages of evolution of the proto-neutron star (PNS), ≤ 10 *ms* after the core bounce at $\rho > 2 \times 10^{14}$ *gcm*⁻³ (see Burrows and Hayes [4] and Epstein [5], and references therein). The evolution of the rapidly rotating PNS during this phase is dominated by vigorous entropy-driven convective motions. Contrary to the arguments given by Janka and Raffelt [6], concerning the existence of a structured PNS (mantle and atmosphere), we suggest that at this post-bounce evolutionary stage there is no room for such a stratified structure to appear. Simply, convection (overturn timescale ~ 1 *ms*) dominates the PNS evolution, what results is a nonx structured PNS at all. Consequently, the analysis of Ref. [6] is not applicable to our model. Instead, we follow the lines of Kusenko and Segrè [3] in what concerns to the oscillation mechanism itself and its implications. Furtherly, since we expect the neutrino oscillation-driven GWs burst to be emitted at core bounce (mass $M_{core} = 1.4 M_\odot$, radius $R \sim 10^7$ *cm* [7], timescale $\tau_B \sim 1$ *ms* [8]), its timescale is a very short one compared to the neutrino diffusion time (the PNS Kelvin-Helmholtz neutrino cooling) ~ 10 *s*, the process itself can be considered as a *transient stage* [6]. As pointed out by Ref. [6] for this phase the mechanism proposed by Kusenko and Segrè [9,3] may properly be at work.

The issue of neutrino oscillations has quite recently been invoked to explain a variety of astrophysical and cosmological concerns. Recent researches in the astrophysics of compact objects allow Kusenko & Segrè [3] and Grasso, Nunokawa & Valle (GNV98) [10] to claim

that the pulsar velocities (recoil kicks) may be explained assuming anisotropic neutrino emission induced by $\nu_{\mu,\tau}$ oscillations into ν_e , and also sterile-to-active oscillations of massive [9,3] or massless [10] neutrinos, provided a strong magnetic field ($B \sim 10^{15}$ G) be involved. Meanwhile, Akhmedov, Lanza and Sciama [10] have studied the influence of resonant spin flavor precession of neutrinos during core-collapse supernova explosion leaving neutron stars as remnants. Their idea is based on the Mikheev, Smirnov & Wolfenstein (MSW) neutrino resonance [11] occurring on a surface off-centric to a proto-neutron star core. The neutrino relative recoil momentum is defined as [9,3]

$$\frac{\Delta\kappa}{\kappa} = \left[\frac{4e\sqrt{2}}{\pi^2} \right] \left(\frac{\mu_e \mu_n^{1/2}}{m_n^{3/2} T^2} \right) B = 0.01 \left(\frac{B}{1.2 \times 10^{15} \text{G}} \right). \quad (1)$$

In this equation μ_e and μ_n , define the electron and neutron chemical potential, respectively, m_n is the neutron mass, and $T \sim 30$ MeV the temperature at the s -neutrinosphere. This anisotropic temperature distribution (the matter-induced neutrino potentials) is created by the strong magnetic field ($B \sim 10^{15}$ G) [9] developed during the supernova core collapse. So, the resonant conversions of ordinary neutrinos (ν_{μ}) into sterile ones (ν_s), if they exist, can occur all the way through the proto-neutron star (PNS) structure involving huge amounts of non-spherically distributed energy. This asymmetric sterile neutrino energy configuration is a crucial point for the arguments presented below supporting the idea that some gravitational-wave (GWs) bursts are expected to be generated by the same time the active-to-sterile neutrino oscillations develop. This ν -oscillation view seems to be confirmed by the Super-Kamiokande Neutrino Experiment (SNE), whose most recent results, published by Fukuda et al. [12], have presented compelling evidence for neutrino oscillations.

Core-collapse supernovae explosions are one of the most powerful sources of neutrinos $\nu_e, \nu_{\mu}, \nu_{\tau}$ and probably ν_s and its antiparticles. Different theoretical and numerical models of type II supernovae explosions [13] have estimated that $\Delta E_{total} = 5.2 \times 10^{53} \text{erg} \left(\frac{10 \text{ km}}{R_{NS}} \right) \left(\frac{M_{NS}}{1.4 M_{\odot}} \right)^2$ are carried away by neutrinos (see below). Almost $\sim 10^{58}$ neutrinos of mean energies $(10 - 25) \text{ MeV}$ are released over time scales of seconds. Investigations have shown that nearly 99% of the total gravitational binding energy of a protoneutron star can directly be carried away by these neutrinos during the supernova explosion [13,14], the remainings streaming away as photons and gravitational waves [14] -a small part of it $\sim (10^{-6} - 10^{-7}) M_{\odot}$ -generated through the quadrupolar distortion of its mass tensor. Because of ν_{μ} and ν_{τ} interact only via neutral currents their neutrino-spheres are deeper in the core, therefore their mean energies are higher, near $(15-25)$

MeV. At this energy level, detection of ν_e and $\bar{\nu}_e$ may occur since they basically undergo charged-current interactions, while as quoted above ν_{μ} and ν_{τ} interact through their neutral-currents [11]. This last point reinforces why their detection is crucial to test the *supernova mechanism*.

In nuclear matter optical depths (mean free paths) are different for different neutrino species. Then they can undergo $\nu_e \longleftrightarrow \nu_{\mu,\tau}$ conversions through the charged-current interactions. Whenever mixing and ν_{τ} mass are non-vanishing, the spherical symmetry of the resonance surface for $\nu_e \longleftrightarrow \nu_{\tau}$ oscillations there doesn't exist any longer, and an asymmetric distribution of the emitted neutrino momenta can occur. The resonance condition for conversions in between neutrinos of left-handed of flavor i into right-handed of flavor j ($i, j = e, \mu, \tau$ or s , with s representing the sterile neutrino) is given by [11,15]

$$V(\nu_{iL}) + \frac{m_{\nu_i}^2}{2E_{\nu}} = V(\bar{\nu}_{jR}) + \frac{m_{\nu_j}^2}{2E_{\nu}}, \quad (2)$$

with $V(\nu_s) = 0$. In the case of magnetized material media it reads:

$$\sqrt{2}G_F N_{eff} - c_{eff} B_{||} = \frac{\Delta m^2}{2E_{\nu}} \cos 2\theta_0. \quad (3)$$

An extensive discussion of neutrino physics including mean potential energies of active neutrinos and antineutrinos is approached to in Ref. [15], and references therein. In the last equation the effective parameters for the resonant conditions, N_{eff} and c_{eff} , depend on the exact nature of the oscillation involved [9]. Since the mass difference constraints the occurrence of a specific resonant conversion, we assume a hierarchical neutrino mass scheme: $\nu_s \gg \nu_{\tau} \gg \nu_{\mu}$. Thus $\nu_s \leftrightarrow \bar{\nu}_{\tau,\mu}$ neutrino oscillations are allowed.

However, Smirnov [15] has called the attention on the possibility that a very strong suppression of the $\nu_{\mu} - \nu_{\tau,s}$ mixing angle can occur in this picture due to the fact that for transitions in extremely dense matter such as the ones prevalent in the supernova at core bounce, $\rho_c \approx (10^{14} - 10^{15}) \text{g/cm}^3$, a rather low mixing angle develops, since the adiabaticity condition should be fulfilled for oscillation lengths

$$L_{\nu_{\mu} \rightarrow \nu_s} \simeq \left([(2\pi)^{-1} \frac{\Delta m^2}{2\kappa}] \sin 2\theta \right)^{-1} \approx \left(\frac{10^{-2}}{\sin 2\theta} \right) \text{cm}, \quad (4)$$

smaller than the typical density variation scale-height ($H_{N_e} \sim 6 \text{ km}$). This constraint is kept if the mixing angle $\sin^2 2\theta > 10^{-8}$. It is easy to check the degree of suppression to be around eight orders of magnitude smaller than the one Super-Kamiokande Experiment has inferred to. However, since during its ploughing through the PNS core ν_{μ} -neutrinos can be absorbed

and reemitted $\sim 10^5$ times, the effective suppression of the $\nu_\mu - \nu_s$ mixing angle reduces to $\sim 10^{-3}$ compare to SKE. This is consistent with the upper limit derived in Ref. [16] for the mass range relevant to our problem: $\Delta m \sim (3-10)$ keV, for which the mixing angle turns out to be: $\sin^2 2\theta_{\bar{e},s} > 10^{-6}$. For larger mixing angle, neutrino detections from SN1987A by IBM and Kamiokande experiments would render contradicted since the electron anti-neutrinos get reduced because most of the PNS energy would be carry off by the sterile neutrino. In the $\bar{\nu}_e \leftrightarrow \nu_s$ case, it is expected that no more than one third of the total neutrino flux is emitted asymmetrically [9]. Moreover, because the explosion produces $\sim 10^{58}$ neutrinos, the minute parcel available can still change flavor forwards and reverse back several times when a radial distance from the proto-neutron star edge equivalent to one mean free path has been reached. There onwards the oscillating sterile neutrinos will carry out information concerning the highly distorted neutrino-sphere at the point of their last conversion. This is the source of the neutrino oscillation-driven gravitational-wave burst.

Simulations of supernovae explosions have demonstrated that the geometry of the neutrino-sphere is not the one of a *real sphere* [14]. High density gradients in the compact object produced as the implosion goes on create anisotropic mass distribution regions where the matter density surpasses the required for neutrino trapping (see [17,13]), while in others they can go through freely. That time-varying anisotropic distribution of density gradients suggest that the energy-momentum tensor of the neutrino fluid is somehow quadrupolar. Then, its time variation translates into the equivalent of a changing quadrupole tensor of mass, whose propagation induces emission of gravitational waves [4]. The other way round, the supernova magnetic field induces the matter content polarization what forces resonant conversions to take place at very radial distances from the star core for those neutrinos released along the magnetic field axis. This yields in a rather inhomogeneous distribution of temperatures for the resonant surface over the dipolar axis, what induces an asymmetric spatial distribution of the radiated neutrino momenta, and with this the possibility that a GWs burst can be triggered.

For the above scenario, we can estimate the characteristics of the gravitational radiation emitted by using Newtonian gravity and the quadrupole approximation to general relativity. Following Burrows and Hayes [4] get

$$h_{ij}^{tt} = \frac{4G}{c^4 D} \int_{-\infty}^t \alpha(t') L_\nu(t') dt'. \quad (5)$$

Here h_{ij}^{tt} represents the transverse-traceless dimensionless metric strain, D defines the distance to the source, $0.2 \leq \alpha(t) \leq 0.8$ [4] is the instantaneous quadrupole anisotropy, $L_\nu(t)$ the overall luminosity of neutrinos. In

our model, we associate the source of the gravitational-wave burst to the quadrupolar moment of the neutrino energy flow, ΔE_ν , by the time the oscillation occurs inside the proto-neutron star, instead of to the PNS quadrupole tensor of mass, as usually done in ordinary neutrino-driven supernova explosion models [14]. With the assumption that the neutrino oscillation timescale is $T_{\nu_\mu \longleftrightarrow \nu_s} \sim c/R_{PNS} \sim 1.0 \times 10^{-3}$ s at the PNS interior, where R_{PNS} is the PNS radius, then we can expect the burst waveform of the radiation emitted to look like a delta Dirac function centred around 1 kHz, superimposed to the overall waveform as computed by Burrows and Hayes [4]. Then the characteristic, normalized, gravitational-wave amplitude generated during the supernova explosion due to the magnetically distorted (non-spherical) outcoming front of the s neutrino-sphere is $h \sim 1.4 \times 10^{-23} \text{ Hz}^{-1/2}$, for a source distance 10 kpc, and in the case for which 100% of the neutrinos undergo oscillations in between $\nu_\mu \longleftrightarrow \nu_{\tau,s}$, with luminosity $L_\nu = 3 \times 10^{54} \text{ erg/s}$ released on a timescale 3×10^{-2} s [5,4]. Unfortunately, a gravitational-wave signal from the Large Magellanic Cloud would not be detectable.

Therefore, a gravitational-wave signal such as this will be detectable by the new generation of gravitational-wave detectors such as LIGO, VIRGO and TIGAs detectors [1,2] provided the large part of neutrino undergo oscillations on a timescale $\sim 1 \text{ ms}$ (see Ref. [8]). The gravitational wave frequency of the main signal (GW burst) produced will be $f_{gw} \sim 1 \text{ kHz}$.

If the mechanism neutrino oscillations/gravitational-wave bursts really realizes in nature, then, it can afford a natural bridge to link and explain two, until now, difficult problems in supernovae physics, i. e., a) what is the very origin of the asymmetries observed in supernovae ejecta and remnants? b) what did trigger the bi-polar beamed jets demonstrated by Nisenson and Papaliolios [18] to be in association with SN 1987A? (see Ref. [19]). It is notheworthy that similar bi-polar jet-like structures have been observed in many other supernovae remnants. For a complete account of this issue see Gaensler [20], and references therein). The general relativistic description of the gravitational radiation generated by escaping neutrinos during a supernova explosion, as demonstrated by Epstein [5], requires that the collapsing star be ellipsoidally deformed due to rotation and anisotropic ν -emission. In this case the metric strain can be expressed by [5]

$$|\theta_{TT}^{ij(\nu)}| \sim 3.6 \times 10^{-21} e^2 \left(\frac{55 \text{ kpc}}{D} \right). \quad (6)$$

where e corresponds to the eccentricity of the ellipsoid projected in the plane through which the oscillating ν s escape. We conjecture here that such an asymmetry e is determined fundamentally by the shape of the neutrino-sphere by the time the oscillation develops. The other way round, the anisotropic neutrino emission responsible,

in the picture by Kusenko and Segrè [3], for the nascent neutron star kicks, might be considered the appropriated scaling of the eccentricity appearing in eq.(5) during supernovae explosions. If it is the case, using Kusenko and Segrè [9,3] results for $B = 10^{15} G$, we can write

$$e^2 \sim \frac{\Delta\kappa}{\kappa} = 0.01, \quad (7)$$

from which we derive the angle θ the final jets will be beamed into driven by neutrino oscillations

$$\tan \theta \simeq e = \frac{a-b}{a+b} = 0.1 \longrightarrow \theta = 5.7 \text{ deg.} \quad (8)$$

This interesting physical connection might provide a simple and elegant explanation to the points raised by Wang and Wheeler [21] concerning the observational features of SN1998bw and other SNe. Based upon spectropolarimetric studies of a large sample of SNe, these authors suggested that sufficiently high energy deposition ($\sim 10^{50} \text{ erg}$) along the polar axis can build oblate isodensity contours in central core collapses, and that the formation of such structures may lead to produce jet-like structures. In our view, the overall process will follow as: the stalled supernova explosion is revived by neutrino-driven convection [14]. Some instants before, the flavor conversions occur, and the large part of ν s is absorbed through β -interactions [3], transferring that way the huge momenta and energy they carry out. The baryon matter in the mantle absorbing the oscillating ν s will keep, and transport outwards, the memory of the asymmetric neutrino absorption. Thus, when these layers finally drive the SNe explosions themselves, they would reflect the aspherical momenta gained at the core. Since, as shown above, the large ν flux is strongly collimated, it is likely then that a strong jetted matter ejecta, and a highly polarized expanding luminous envelope appears. An additional bonus given by the present scenario is the fact the model naturally explains why do the expected emitted jets are to have markedly different strengths, a point also discussed by Cen [19] in his unified picture for pulsar, gamma-ray bursts and supernovae. In our picture, this power asymmetry of jets is fundamentally created by the neutrino oscillation mechanism.

To conclude, we have propounded for the first time that a very fundamental new class of gravitational wave bursts can be produced during resonant *ordinary-to-sterile* neutrino oscillations ($\nu_\mu \leftrightarrow \nu_{\tau,s}$) inside proto-neutron stars at the onset of core-collapse explosions of massive stars. This new family of GWs signals should be viewed as one of a very different nature when compared to the classical quadrupolar mass-tensor gravitational waves, as currently is suggested in studies on this subject, when ordinary neutrinos drive the final (stalled) explosion. The characteristic properties of the signal are found to be in the sensitivity range of most of the now under construc-

tion or in phase of planning gravitational-wave observatories.

This mechanism for GWs emission was not envisaged in former investigations on the subject perhaps because there was no compelling evidence for the neutrino flavor changes. This letter introduces it for the first time, in the context of mechanisms for generating gravitational radiation from astrophysical sources. The eventual detection of such bursts in future experiments will be a breakthrough in fundamental physics for, as pointed out above, there exist the chance that an important parcel of the oscillating sterile neutrinos can get the Earth nearly by the time the associated gravitational radiation bursts they produced when changing flavors ($\nu_\mu \leftrightarrow \nu_{\tau,s}$) arrive at the detectors.

ACKNOWLEDGMENTS

I am grateful to Dr. A. Kusenko and Professor A. Yu. Smirnov for their kindly reading of this manuscript and valuable key suggestions.

-
- [1] K. S. Thorne, *Proceedings of the Snowmass Summer Study on Particle and Nuclear Astrophysics and Cosmology*, eds. E. W. Kolb and R. Peccei, Wprld Scientific, Singapore (1995). *ibid Three Hundred Years of Gravitation*, Eds. S. W. Hawking and W. Israel, Cambridge University Press, Cambridge, England (1987).
 - [2] B.F. Schutz, in *Proceedings of the Les Houches School on Astrophysical Sources of Gravitational Radiation*, eds. J.-P. Lasota and J.-A. Marck, Cambridge University Press, Cambridge, England (1995). B.F. Schutz, report astro-ph/9802020, February (1998).
 - [3] A. Kusenko and G. Segrè, Phys. Rev. Lett. **79**, 2751 (1997). *ibid.* **77**, 4872 (1996), Phys. Rev. D, **59**, 061302 (1999).
 - [4] A. Burrows and J. Hayes, Phys. Rev. Lett. **76**, 352 (1996). See also A. Burrows, J. Hayes and B. A. Fryxell, Astrophys. J. **445**, 830 (1995).
 - [5] R. Epstein, Astrophys. J., **223**, 1037 (1978).
 - [6] H.Th. Janka and G. G. Raffelt, Phys. Rev. D, **59**, 023005 (1999).
 - [7] J. R. Wilson, Astrophys. J., GREP Conference 79 (1977).
 - [8] M. Turner and R. V. Wagoner, Stanford University IT, preprint No. 576 (1977).
 - [9] A. Kusenko and G. Segrè, Phys. Letts. **B396**, 197 (1997).
 - [10] D. Grasso, H. Nunokawa and J.W.F. Valle, Phys. Rev. Lett. **81**, 2412 (1998). See also E.Kh. Akhmedov, A. Lanza and D.W. Sciama, Phys. Rev. D **56**, 6117 (1997). H. Nunokawa, V.B. Semikoz, A.Yu. Smirnov and J.W.F. Valle, Nucl. Phys. **B501**, 17 (1997).
 - [11] S.P. Mikheyev and A.Yu. Smirnov, Yad. Fiz. **42**, 1441 (1985). [Sov. J. Nucl. Phys. **42**, 913 (1985)]; Prog. Part.

- Nucl. Phys. **23**, 41 (1989). L. Wolfenstein, Phys. Rev. D. **17**, 2369 (1978); **20**, 2634 (1979).
- [12] Fukuda, Y., et al., Phys. Rev. Lett. **81**, 1562 (1998).
 - [13] S.E. Woosley and T.A. Weaver, Astrophys. J. S., **101**, 181 (1995). A. Burrows, AAS, **187**, 1704 (1995). *ibid.* **187**, 1703 (1995).
 - [14] H.-Th. Janka and E. Müller, A & A **306**, 167 (1996). E. Müller and H.-Th. Janka, A & A **317**, 140 (1997).
 - [15] Private communication with Professor A. Yu. Smirnov, International Centre for Theoretical Physics (ICTP). See also J.F. Beacom, *Supernova Neutrinos and the Neutrino Masses*, report hep-ph/9901300, January (1999).
 - [16] K. Kainulainen, J. Maalampi and J. T. Peltoniemi, Nuc. Phys. B, **358**, 435 (1991).
 - [17] H.V. Klapdor-Kleingrothaus and K. Zuber, *Particle Astrophysics*, Chapter 13, Institute of Physics Publishing Ltd., Bristol and Philadelphia, USA (1997).
 - [18] P. Nisenson and C. Papaliolios, Report astro-ph/9904109, April 9 (1999).
 - [19] R. Cen, Astrophys. J., **507**, L131 (1998) and Report astro-ph/9904147, April 12 (1999).
 - [20] B. M. Gaensler, Ph. D. Thesis Dissertation, University of Sidney, Australia. Unpublished (1999).
 - [21] L. Wang and J. C. Wheeler, Astrophys. J., **504**, L87 (1998). See also J.C. Wheeler, P. Hoefflich and L. Wang, AAS DPS meeting # 193, # 42.06 (1998), Report astro-ph/9904047, April 3 (1999).